



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/38, 31/40, 31/445, 31/535, 31/55, C07D 333/58, 409/10, 409/12, 413/10, 413/12	A1	(11) International Publication Number: WO 98/48792 (43) International Publication Date: 5 November 1998 (05.11.98)
(21) International Application Number: PCT/US98/08477 (22) International Filing Date: 28 April 1998 (28.04.98) (30) Priority Data: 60/045,132 30 April 1997 (30.04.97) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indi- anapolis, IN 46285 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): MCGILL, John, McNeil, III [US/US]; 1120 Hasp Way, Lafayette, IN 47905 (US). MILLER, Randal, Scot [US/US]; 3814 Constitution Drive, Carmel, IN 46032 (US). (74) Agents: STRODE, Janelle, D. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: A REGIOSELECTIVE ALKYLATION PROCESS FOR PREPARING SUBSTITUTED BENZO[b]THIOPHENES (57) Abstract The present invention provides processes for the regioselective alkylation of benzothiophenes.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**A Regioselective Alkylation Process for Preparing
Substituted Benzo[b]thiophenes**

The present invention relates to chemical processes.
5 More particularly, the present invention relates to a regioselective alkylation process for preparing 2-(4-hydroxyphenyl)-3-[4-(W-substituted amino)alkoxybenzoyl]benzo[b]thiophene compounds.

The compound 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-
10 piperidinoethoxy)benzoyl]benzo[b]thiophene, known also by its generic chemical name, raloxifene, is an important member of the class of compounds known as selective estrogen receptor modulators (SERMs). That compound is disclosed in U.S. Pat. No. 4,418,068.

15 Some of the compounds prepared by the present processes, including raloxifene, were first described in U.S. Pat. No. 4,133,814. That patent taught the use of phenacyl, halophenacyl, and alkyl protecting groups for phenolic hydroxyl groups. Those alkyl protecting groups may
20 be removed by treating the phenolic ethers with pyridine hydrochloride.


The process described in U.S. Pat. No. 4,358,593 used additional protecting groups (for example, acetyl, substituted acetyl, benzoyl, alkylsulfonyl, and arylsulfonyl
25 groups) for preparing 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-aminoethoxy)benzoyl]-benzo[b]thiophenes. That patent further taught the use of classical Friedel-Crafts catalysts in the acylation of the protected 2-(4-hydroxyphenyl)-6-hydroxybenzo[b]thiophene, including metal halides such as
30 aluminum chloride, aluminum bromide, zinc chloride, boron trifluoride, boron tribromide, titanium tetrachloride, titanium tetrabromide, stannic chloride, stannic bromide, bismuth trichloride, and ferric chloride. Subsequent to

- 2 -

acylation, the protecting group was generally removed under basic conditions.


The process described in U.S. Pat. No. 5,470,854 taught the use of methyl protecting groups that could be cleaved in a one pot process during the Friedel-Crafts acylation by particular acylation catalysts. That patent taught that boron halides, such as boron trichloride and boron tribromide, were particularly useful in the Friedel-Crafts acylation, and for the cleavage of arylmethyl ethers.

The present invention provides a process for preparing compounds of formula VI



VI


wherein R^1 and R^2 each are independently C_1 - C_4 alkyl, or combine to form, together with the nitrogen to which they are attached, a piperidinyl, pyrrolidinyl, methylpyrrolidinyl, dimethylpyrrolidinyl, morpholino, dimethylamino, diethylamino, or 1-hexamethyleneimino ring, and n is 2 or 3, which includes the reaction of a compound of formula IV



IV

- 3 -

with a compound of formula V



V

5

wherein Y is chloro or 4-methylphenylsulfonyl-O-, and R¹ and R² and n are as defined above, in the presence of a suitable base.

General terms used in the description of chemical
10 formula herein bear their usual meanings. For example, the term "C₁-C₄ alkyl" refers to straight or branched chains of 1 to 4 carbon atoms including, methyl, ethyl, propyl, isopropyl, butyl, n-butyl, and isobutyl; and the term "alkyl" encompasses the groups included in the definition of
15 "C₁-C₄ alkyl" in addition to groups such as pentyl, isopentyl, hexyl, isohexyl, and the like. The term "halo" includes bromo, chloro, fluoro, and iodo. The term "lower alcohols" refers to C₁-C₄ alcohols including methanol, ethanol, propanol, isopropanol, butanol, n-butanol,
20 isobutanol, and the like. The abbreviation "Tos" refers to a 4-methylphenylsulfonyl group.

The term "chlorinating reagent" encompasses such reagents as thionyl chloride, molecular chlorine, N-chlorosuccinimide, benzeneseleninylchloride/aluminum
25 chloride, and the like.

The term "hydroxy protecting group" denotes a group understood by one skilled in the organic chemical arts of the type described in Chapter 2 of "Protective Groups in Organic Synthesis, 2nd Edition, T. H. Greene, et al., John
30 Wiley & Sons, New York, 1991. Such groups include, for example, ether groups, including methyl and substituted methyl ether groups such as methyl ether, methoxymethyl

- 4 -

ether, methylthiomethyl ether, *tert*-butylthiomethyl ether, (phenyldimethylsilyl)methoxymethyl ether, benzyloxymethyl ether, *p*-methoxy-benzyloxymethyl ether, and *tert*-butoxymethyl ether; substituted ethyl ether groups such as
5 ethoxyethyl ether, 1-(2-chloroethoxy)ethyl ether, 2,2,2-trichloroethoxymethyl ether, and 2-(trimethylsilyl)ethyl ether; phenyl and substituted phenyl ether groups such as phenyl ether, *p*-chlorophenyl ether, *p*-methoxyphenyl ether, and 2,4-dinitrophenyl ether; benzyl and substituted benzyl
10 ether groups such as benzyl ether, *p*-methoxybenzyl ether, *o*-nitrobenzyl ether, and 2,6-dichlorobenzyl ether; and alkylsilyl ether groups such as trimethyl- triethyl- and triisopropylsilyl ethers, mixed alkylsilyl ether groups such as dimethylisopropylsilyl ether, and diethylisopropylsilyl
15 ether; and ester protecting groups such as formate ester, benzylformate ester, mono- di- and trichloroacetate esters, phenoxyacetate ester, and *p*-chlorophenoxyacetate and the like. Preferred protecting groups encompassed in this invention are methyl groups, for example wherein R is
20 methyl.

Methods for protecting hydroxy groups with one of the groups listed above, or as provided in the text by Greene, *et al.*, as well as methods for cleaving or removing the protecting groups, are discussed in the text and the
25 references cited therein and, are well known to one skilled in the art.

The term "acyl activating group" refers to a substituent to a carbonyl that promotes nucleophilic addition reactions to the carbonyl. Suitable activating
30 substituents are those which have a net electron withdrawing effect on the carbonyl. Typical electron withdrawing groups include groups that when combined with the carbonyl form an ester or amide. Such groups include hydroxybenzotriazole, imidazole, nitrophenol, pentachlorophenol, N-

- 5 -

hydroxysuccinimide, dicyclohexylcarbodiimide, N-hydroxy-N-methoxyamine, and the like. The term acyl activating group also encompasses groups that when combined with the carbonyl form an acid anhydride. Such groups include small

5 carboxylic acids such as acetic, formic, sulfonic, methanesulfonic, ethanesulfonic, benzenesulfonic, or p-tolylsulfonic acid, and the like. Furthermore, a halogen attached to a carbonyl activates it for nucleophilic addition. Suitable halogens include chloro, bromo, or iodo.


10 By the term "Lewis acid catalyst" is meant catalysts including metal halides, such as aluminum bromide, aluminum chloride, boron trifluoride, boron trichloride, boron tribromide, titanium tetrachloride, titanium tetrabromide, stannic chloride, stannic bromide, bismuth trichloride, and
15 ferric chloride.

A reaction which is run "neat" is a reaction where one of the reagents also serves as a solvent.

The term "kinetic base" refers to a base which provides a non-reversible deprotonation of an acidic substrate, such
20 as alkyl metals (such as butyl lithium or ethyl magnesium bromide), metal amides (such as lithium diisopropyl amide), or a metal hydride (such as sodium hydride, lithium hydride).

The overall process of the present invention, including
25 the preparation of starting materials, is depicted in Scheme 1 wherein R is independently at each occurrence a hydroxy protecting group, X is an acyl activating group, and n, R¹, R², and Y are as described, *supra*.

- 6 -

Scheme 1

5 A compound of formula I may be acylated with a compound
 of formula II. For example, a compound of formula I may be
 dissolved in a suitable solvent in the presence of Lewis
 acid catalyst and a compound of formula II added. Suitable
 solvents include chlorinated alkane solvents, such as
 10 chloroform, 1,2-dichloroethane, 1,2,3-trichloropropane,
 aromatic solvents such as benzene and chlorobenzene,
 hydrocarbon solvents such as petroleum ether, hexane,
 mixtures thereof, and the like. Methylene chloride is
 typically the preferred solvent. Suitable Lewis acid
 15 catalysts are of the type described, *supra*, or in Olah,
 "Friedel-Crafts and Related Reactions," Interscience
 Publishing Co., New York, 1963. Aluminum chloride or boron
 trichloride is the preferred Lewis acid catalyst. The Lewis
 acid catalyst is typically employed in a substantial molar
 20 excess. For example, a 1.5 to 2.5 molar excess, relative to
 the compound of formula I, is generally employed. A 2.0
 molar excess is preferred. The compound of formula II is
 typically employed in a slight molar excess. For example, a

- 7 -

1.01 to 1.25 molar excess, relative to the compound of formula I is generally employed. A 1.1 molar excess is preferred. The reaction is preferably carried out at about room temperature for about 1 hour.

5 When boron trichloride or boron tribromide is the Lewis acid catalyst employed, the hydroxy protecting group at the 3 position of the benzothiophene ring in compounds of formula III is not removed in the acylation step. The removal of the hydroxy protecting groups at the 2 and 6
10 positions is not important in the overall process because in the next step all of the protecting groups are deliberately removed.

 A compound of formula IV may be obtained by treating a compound of formula III in a suitable solvent with
15 pyridinium hydrochloride. Generally, and preferably, the reaction is run neat in pyridinium hydrochloride. The pyridinium hydrochloride is typically employed in a substantial molar excess. For example, a 15 to 30 molar excess, relative to the compound of formula III is generally
20 employed. A 20 molar excess is preferred. The reaction is preferably carried out at about 185 °C for about 1.5 hours.

 A compound of formula VI may be obtained by the dissolution or suspension of a compound of formula IV in a suitable polar solvent in the presence of a suitable base
25 followed by the addition of a compound of formula V. Suitable solvents include dimethylformamide, lower alcohols, 1,3-dimethyl-2-imidazolidinone, mixtures thereof, and the like. Dimethylsulfoxide is the preferred solvent. Suitable bases include non-kinetic bases such as carbonates,
30 bicarbonates, or hydroxides, (for example, sodium hydroxide, potassium bicarbonate, or sodium hydroxide) and kinetic bases such as alkyl lithiums (for example, n-butyl lithium) or sodium hydride. A kinetic base, specifically sodium hydride, is the preferred base. The base is typically

- 8 -


employed in a slight molar excess. For example, a 1.01 to 1.15 molar excess, relative to the compound of formula IV, is generally employed. A 1.05 molar excess is preferred. A compound of formula V is typically employed in a slight molar excess. For example, a 1.01 to 1.15 molar excess, relative to the compound of formula IV is generally employed. A 1.05 molar excess is preferred. The reaction is preferably carried out at about 35 °C for about 2 hours after addition of the base and then at about 65 °C for about 16 hours after the addition of a compound of formula V.

Surprisingly, when the alkylation is performed as described in this invention, the alkylation proceeds in a regioselective fashion to give a compound of formula VI.

Compounds of formula I are known in the art and are prepared, for example, as described by Peters in U.S. Pat. No. 4,380,635, or Jones, et al., in U.S. Pat. Nos. 4,133,814 and 4,418,068. Compounds of formula II are known in the art and are generally commercially available or prepared by methods well known in the art from readily available starting materials.

Compounds of formula V are commercially available or may be prepared from commercially available compounds of formula VII as shown in Scheme 2 wherein n, R¹, and R² are as defined, *supra*.

- 9 -

Scheme 2

5 Compounds of formula V include compounds of both
 formula Va and compounds of formula Vb as illustrated above.
 For example, compounds of formula V where Y is chloro may be
 prepared from compounds of formula VII by dissolving a
 compound of formula VII in a suitable solvent and adding
 10 thionyl chloride or 2,2,2-trifluoroethylsulfonyl chloride.
 Similarly, compounds of formula V where Y is 4-
 methylphenylsulfonyl-O- may be prepared from compounds of
 formula VII by the addition of 4-methylphenylsulfonyl
 chloride. Suitable solvents include acetonitrile,
 15 chloroform, ethyl acetate, dimethylformamide,
 tetrahydrofuran, lower alcohols, and the like. A preferred
 solvent is methylene chloride. The additive is typically
 employed in a slight molar excess. For example, a 1.01 to
 1.3 molar excess, relative to a compound of formula VII, is
 20 generally employed. A 1.12 molar excess is typically
 preferred. The reaction is typically and preferably carried
 out at about 0 °C during the addition and then at about room
 temperature for about 12 hours. A preferred embodiment of
 this invention employs 2-chloroethylpiperidine (a compound
 25 of formula V wherein n is 2, R¹ and R² combine to form
 piperidiny1, and Y is chloro) generated and used *in situ*

- 10 -

from 2-hydroxyethylpiperidine (a compound of formula VII where n is 2 and R¹ and R² combine to form piperidinyl) by the process described in Scheme 2, in the overall process of the invention described in Scheme 1.

5 In general, the reactions of Schemes 1 and 2 are substantially complete in about 15 minutes to 72 hours when conducted at a temperature range of from about 0 °C to the reflux temperature of the reaction mixture. The reaction solvent choices are not critical so long as the solvent
10 employed is inert to the ongoing reaction and the reactants are sufficiently solubilized to effect the desired reaction. Once a reaction is complete, the intermediate compound may be isolated, if desired, by procedures known in the art. For example, the compound may be crystallized and then
15 collected by filtration, or the reaction solvent may be removed by extraction, evaporation, or decantation. The intermediate may be further purified, if desired by common techniques such as recrystallization or chromatography over solid supports such as silica gel or alumina. The compounds
20 of formula I are preferably isolated before use in subsequent reactions.

The optimal time for performing the reactions of the invention can be determined by monitoring the progress of the reaction via conventional chromatographic techniques.
25 Furthermore, it is preferred to conduct the reactions of the invention under an inert atmosphere, such as, for example, argon or nitrogen.

The following examples are provided for the purpose of illustrating the present invention and are not intended to
30 be limiting upon the scope of the invention.

- 11 -

ExamplesPreparation 15 2-(4-Methoxyphenyl)-3-(4-Methoxybenzoyl)-6-
 Methoxybenzo[b]thiophene

A 250 mL, 3 neck, round bottomed flask fitted with a mechanical agitator and connected to a sodium hydroxide
10 scrubber is purged with nitrogen and charged with 2-(4-methoxyphenyl)-6-methoxybenzo[b]thiophene (5.3 g, 19.6 mmol), p-anisoyl chloride (3.68 g, 21.56 mmol), and 100 mL of methylene chloride. To this slurry at 20 °C is added boron trichloride (3.4 mL, 39.8 mmol) which had been
15 condensed in a graduated cylinder. The reaction mixture is quenched after 40 minutes by the slow addition of 50 mL of methanol over approximately 5 minutes followed by addition of 50 mL of water over 5 minutes. The organic phase is washed with 50 mL brine, dried over sodium sulfate,
20 filtered, and concentrated affording a dark orange oil. Addition of 10 mL of methanol followed by sonication for 5 minutes resulted in crystallization of 7.1 g of crude product as a yellow solid. The crude material is recrystallized from 3:1 hexanes/toluene affording 5.9 g of
25 product as pale yellow crystals. (75%).
mp 118-119 °C; ¹H NMR (300.1 MHz, DMSO-d₆) δ 3.66 (s, 3H), 3.77 (s, 3H), 3.86 (s, 3H), 6.84 (d, 2H), 6.88 (d, 2H), 6.98 (dd, 1H), 7.32 (d, 2H), 7.34 (s, 1H), 7.63 (d, 1H), 7.68 (d, 2H); ¹³C NMR (75.5 MHz, DMSO-d₆) δ 55.2, 55.5, 105.2, 114.1, 114.4, 115.0, 123.3, 125.2, 129.6, 129.7, 130.2, 131.8, 133.2, 139.4, 140.8, 157.4, 159.5, 163.6, 192.4; MS (FD+) m/e 405 (30), 404 (100); Analysis calc'd for C₂₄H₂₀O₄S: C, 71.27; H, 4.98; S, 7.93; found: C, 71.51; H, 4.99; S, 7.98.

- 12 -

Preparation 22-(4-Hydroxyphenyl)-3-(4-Methoxybenzoyl)-6-
Hydroxybenzo[b]thiophene

5

A 100 mL, 3 neck, round bottom flask fitted with a mechanical agitator, glass stopper, and outlet to a caustic scrubber is charged with p-anisoyl chloride (3.47 g, 20.34 mmol), 2-(4-methoxyphenyl)-3-(4-methoxybenzoyl)-6-methoxybenzo[b]thiophene (5.0 g, 18.49 mmol) and 30 mL 1,2-dichloroethane. The mixture is cooled to 0-5°C and previously condensed boron trichloride (4.7 mL, 55.0 mmol) is added in one portion. The mixture is stirred at 0-5°C for 7.5 hours then more boron trichloride (4.7 mL, 55.0 mmol) is added. The cooling bath is removed and the reaction mixture is stirred at ambient temperature for 16 hours. After the overnight stir, the mixture is poured into 500 mL of agitated ice/water which caused precipitation of the monomethoxy product. The reaction vessel is rinsed in with 1,2-dichloroethane and the slurry is stirred for 1 hour. Then, 200 mL of methylene chloride is added and the mixture is vacuum filtered washing with methylene chloride. The solid is dried overnight in a vacuum oven at 40°C to afford 6.3 g (82.4%) of monomethoxy product as a yellow solid. An analytical sample is obtained by chromatography on silica gel (20:1 methylene chloride / methanol). mp 127-128 °C (dec); ¹H NMR (300.1 MHz, DMSO-d₆) δ 3.76 (s, 3H), 6.66 (d, 2H), 6.85 (dd, 1H), 6.87 (d, 2H), 7.16 (d, 2H), 7.25 (d, 1H), 7.33 (s, 1H), 7.66 (d, 2H), 9.76 (br s, 2H); ¹³C NMR (75.5 MHz, DMSO-d₆) δ 55.5, 107.1, 114.0, 115.2, 115.7, 123.3, 123.8, 129.7, 131.8, 132.3, 139.2, 140.4, 155.4, 157.8, 163.5, 192.6; MS (FD+) m/e 376(100), 377(25), 378(7); Analysis calc'd for C₂₂H₁₆O₄S: C, 70.20; H, 4.28; S, 8.52; found: C, 70.09; H, 4.27; S, 8.45.

- 13 -

Preparation 32-Chloroethylpiperidine

5

Into a 50 mL round bottom flask is placed 6.40 g (33.6 mmol) of *p*-toluenesulfonyl chloride and 25 mL of methylene chloride. The resulting solution is cooled with an ice bath as 4.00 g (31.0 mmol) of 1-piperidinoethanol in 6 mL of methylene chloride is added dropwise. After the addition is complete, the ice bath is removed and the resulting slurry is stirred for about 12 hours. The reaction mixture is concentrated on a rotary evaporator to yield a solid residue which may then be used as in Preparation 4 below.

15

Preparation 42-(4-Hydroxyphenyl)-3-(4-Hydroxybenzoyl)-6-Hydroxybenzo[b]thiophene

20 A 250 mL, 3 neck, flask fitted with a thermometer, condenser, and stopper is charged 2-(4-hydroxyphenyl)-3-(4-methoxybenzoyl)-6-hydroxybenzo[b]thiophene (3.0 g, 7.97 mmol) and pyridine hydrochloride (18.5 g, 160 mmol). The contents are heated to 185-190 °C and held at this temperature for 90 minutes. The hot mixture is then poured into 250 mL of ice/water. A yellow solid formed with this addition. The material is extracted into 250 mL of ethyl acetate. The aqueous is then back-extracted with 100 mL of ethyl acetate. The combined organics are washed with 150 mL 30 1N hydrochloric acid and 150 mL brine. The solvent is removed *in vacuo* giving a viscous oil. To the oil is added 10 mL of 1:1 methylene chloride/hexanes which induced crystallization. The slurry is cooled to 0 °C and stirred at this temperature for 1 hour. The product is collected by

- 14 -

vacuum filtration and washed with 25 mL of 4:1 hexanes/methylene chloride. Drying the solid at 100 °C for 5 hours afforded 2.5 g (86.2%) of a dark yellow solid containing approximately 3% monomethoxy starting material.

5 An analytical sample is obtained by chromatography on silica gel (15:1 methylene chloride:methanol). mp 135 °C (dec); ¹H NMR (300.1 MHz, DMSO-d₆) δ 6.67 (d, 2H), 6.71 (d, 2H), 6.85 (dd, 1H), 7.17 (d, 2H), 7.25 (d, 1H), 7.32 (d, 1H), 7.57 (d, 2H), 9.73 (br s, 2H), 10.41 (br s, 1H); ¹³C NMR (75.5 MHz, DMSO-d₆) δ 107.1, 115.1, 115.4, 115.6, 123.4, 123.9, 128.4, 129.7, 129.9, 132.1, 132.4, 139.2, 140.0, 155.4, 157.8, 162.5, 192.4; MS (FD+) m/e 362 (100), 363 (24), 364 (7); HRMS m/e calc'd for C₂₁H₁₅O₄S (M+1): 363.069106, found 363.06990.

15

Example 1

[6-Hydroxy-2-(4-Hydroxyphenyl)-3-[4-(2-Piperidinoethoxy)benzyl]benzo[b]thiophene Hydrochloride]

20 A 50 mL, 3 neck, round bottomed flask fitted with a mechanical agitator, nitrogen purging inlet, and thermometer is charged with 2-(4-hydroxyphenyl)-3-(4-hydroxybenzoyl)-6-hydroxybenzo[b]thiophene (1.0 g, 2.76 mmol) and 25 mL dimethylsulfoxide. The solution is then charged with sodium
25 hydride (60% oil dispersion, 116 mg, 2.90 mmol) in one portion. A large amount of gas evolution is noted with this addition. The temperature of the mixture is adjusted to 35 °C and the reaction allowed to stir at this temperature for 2 hours. 2-chloroethylpiperidine (428 mg, 2.90 mmol) is
30 then added via syringe in one portion. The 2-chloroethylpiperidine is rinsed with 0.3 mL dimethylsulfoxide. The reaction mixture is then warmed to 65 °C and stirred for 16 hours. The mixture is then quenched by the slow addition of 25 mL deionized water. The


- 15 -

contents are transferred to a 250 mL Erlenmeyer flask and another 50 mL deionized water is added. After stirring vigorously for 1 hour, the solid is collected by vacuum filtration and washed with water. After drying, 1.3 g of a mixture of 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinoethoxy)benzyl]benzo[b]thiophene and trihydroxy starting material is obtained. Purification by flash chromatography (methanol) afforded 550 mg of a viscous yellow oil which is crystallized from 20:1 methanol/ water at pH 1 giving 370 mg of a light yellow powder. mp 252-256 °C (Lit. 258°C); ¹H NMR (300.1 MHz, DMSO-d₆) δ 1.32 (m, 1H), 1.65 (m, 1H), 1.73 (br s, 4H), 2.93 (m, 2H), 3.40 (m, 4H), 3.62 (br s, 1H), 4.45 (m, 2H), 6.68 (d, 2H), 6.83 (dd, 1H), 6.94 (d, 2H), 7.13 (d, 2H), 7.24 (d, 1H), 7.37 (d, 1H), 7.58 (d, 2H), 9.88 (br s, 1H), 10.46 (br s, 1H); ¹³C NMR (75.5 MHz, DMSO-d₆) δ 23.9, 25.5, 25.6, 54.3, 54.4, 57.1, 65.9, 107.1, 114.5, 115.2, 115.7, 123.3, 123.6, 129.63, 129.64, 129.7, 131.8, 132.2, 139.2, 140.3, 155.6, 158.0, 162.8, 192.6; MS (FAB+) m/e 474 (91), 155 (70), 152 (97), 135 (99), 119 (100); Analysis calc'd for C₂₈H₂₈NO₄SCl: C, 65.99; H, 5.54; N, 2.75; Cl, 6.87; S, 6.28; found: C, 64.78; H, 5.49; N, 2.86; Cl, 7.04; S, 6.33.

- 16 -

We claim

1. A process for preparing a compound of formula VI



VI


wherein:

R^1 and R^2 each are independently C_1 - C_4 alkyl, or
 10 combine to form, together with the nitrogen atom to which
 they are attached, a piperidinyl, pyrrolidinyl,
 methylpyrrolidinyl, dimethylpyrrolidinyl, morpholino,
 dimethylamino, diethylamino, or 1-hexamethyleneimino ring;
 and

15 n is 2 or 3;


which comprises:

reacting a compound of formula IV



IV

with a compound of formula V



V


- 17 -

wherein Y is chloro or 4-methylphenylsulfonyl-O-, and n, R¹, and R² are as defined above;
in the presence of a suitable base.

5

2. A process according to Claim 1 wherein said suitable base is sodium hydride.

3. A process according to Claim 1 wherein said
10 compound of formula V is generated from the reaction of a chlorinating reagent and a compound of formula VII



VII

15

and the resulting compound of formula V is employed *in situ*.

20

4. A process according to Claim 2 wherein said chlorinating reagent is thionyl chloride.

5. A process according to Claim 1 wherein n is 2 and R¹ and R² combine to form, together with the nitrogen to which they are attached, a piperidinyl ring.

25

6. A process according to Claim 1 wherein Y is chloro.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/08477

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US-CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS-structure, Dialog--chem. literature

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,472,962 A (KOIZUMI ET AL) 05 December 1995, see entire document, especially column 4-5.	1-66
Y	US 3,983,245 A (LADD ET AL) 28 September 1976, see entire document, especially column 3-4, schem C.	1-6
Y	US 4,007,204 A (DESCAMPS ET AL) 08 February 1977, see entire document, especially column 2.	1-6



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

19 AUGUST 1998

Date of mailing of the international search report

25 September 1998 (25.09.98)

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized Officer

CELIA CHANG

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/08477

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	JONES ET AL Antiestrogens. 2. Structure-activity studies in a series of 3-aryl-2-arylbenzo [b] thiophene derivatives leading to [6-hydroxy-2 (4-hydroxyphenyl) benzo [b] thie-3-yl] [4- [2- (1-piperidinyl) ethoxy] - phenyl] methanone hydrochloride (LY156758), a remarkable effective estrogen antagonist with only minimal intrinsic estrogenicity. J. Med. Chem. 1984, Vol. 27, pages 1057-1064. see the whole article, especially page 1059.	1-66
Y	WO 89/02893 A1 (THE UNIVERSITY OF TENNESSEE RESEARCH CORPORATION) 06 April 1989, see entire document, especially page 4.	1-66
Y	WO 95/10513 A1 (PFIZER INC.) 20 April 1995, see entire document, especially pages 15 and 39, formula 5a-1 and reaction scheme.	1-6

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/08477

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 31/38, 31/40, 31/445, 31/535, 31/55; C07D 333/58, 409/10, 409/12, 413/10, 413/12

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/212, 233.5, 324, 422, 448; 540/596; 544/145; 546/202; 548/525, 527; 549/51; 564/503, 510

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

514/212, 233.5, 324, 422, 448; 540/596; 544/145; 546/202; 548/525, 527; 549/51; 564/73, 123; 568/74, 77, 306